

## The identity of *Testudo punctata* Lacepède, 1788 (Testudines, Trionychidae)

by Robert G. WEBB \*

**Résumé.** — Après la redescription de l'holotype de *Testudo punctata* Lacepède, 1788, l'histoire taxonomique de cette espèce est débattue ; pour la première fois *Trionyx coromandelicus* Geoffroy, 1809, est considéré comme un synonyme objectif de celle-ci, étant décrit d'après le même spécimen. De plus il apparaît que cet holotype est très probablement originaire de l'Inde péninsulaire : à la sous-espèce nominale *Lissemys punctata punctata* (Lacepède, 1788) correspond donc la forme habituellement désignée *Lissemys punctata granosa* (Schoepff, 1792), et un nouveau nom est proposé pour la sous-espèce taehetée particulière à la plaine indo-gangétique.

**Abstract.** — The holotype of *Testudo punctata* Lacepède, 1788, is redescribed, and the nomenclatural and taxonomic history of that name is discussed. Novel conclusions are the synonymy of *Trionyx coromandelicus* Geoffroy, 1809, and *T. punctata* (objective synonyms, based on same specimen), and determination of the geographical provenance of the holotype of *T. punctata* as most probably peninsular India. The nominal subspecies *Lissemys punctata punctata* is regarded as the recognizable taxon in peninsular India and Sri Lanka (currently referred to as *Lissemys punctata granosa*). Strangely, there is no available name for the common, yellow-spotted, Indus-Gangetic *Lissemys*, and a new name is proposed.

---

### DESCRIPTION OF *Testudo punctata* Lacepède

Following SMITH (1931 : 157), some subsequent authors have credited the name *Testudo punctata* to BONNATERRE (1789 : 30) as the first author to use binomial nomenclature. Although LACEPÈDE referred to his new species only as "La Chagrinée" in the descriptive remarks (1788 : 171), and in the legend of the illustration (Pl. XI), he did apply binomial nomenclature, "*T. punctata*", on the second of two folded charts entitled "Synopsis methodica Quadrupedem oviparorum". The name "T[ortue] chagrinée" appears on the first folded chart, "Table méthodique des Quadrupèdes ovipares"; all names listed there are in the context of common names. WERMUTH (1956 : 409) previously discussed the priority of LACEPÈDE over BONNATERRE, as did SAVAGE (1952 : 204) in his discussion of LACEPÈDE's *Coluber boiga*. LACEPÈDE's turtles work (with folded charts) is quarto size and included in Tome 1; his complete work has been printed in two separate volumes, and the "charts" in French and Latin have been disposed of either as French in Tome 1 and Latin in Tome 2 (published 1789), or vice versa, or the charts are absent (Roger BOUR, pers. comm.).

LACEPÈDE's description of "La Chagrinée" (1788 : 171-172) from "des grandes Indes au Cabinet du Roi, par M. Sonnerat", is based on one specimen; he gave measurements,

\* Department of Biological Sciences, The University of Texas at El Paso, El Paso, Texas 79968, USA.

and mentioned eight pairs of costals and six neurals on the carapace and seven plastral callosities. LACEPÈDE did not mention any features of pattern. LACEPÈDE did mention the name "Chagrinée" (and presumably the specific name, *punctata*) is in allusion to "de petits points" on the callosities of the carapace. The prominence of the bony carapace (callosities and individual costals and neurals) suggests that the holotype at the time of LACEPÈDE's description was not alive and was a dried specimen (probably for some time). LACEPÈDE's illustration of the holotype (Pl. XI) has since been copied by BONNATERRE (1789, Pl. 6, Fig. 4), SCHNEIDER (1789, Pl. unnumbered), DAUDIN (1801, Pl. XIX, Fig. 2), and by SHAW (1802, Pl. 14, upper) all of whom also provided some descriptive remarks.

The holotype of *Testudo punctata* Lacepède is still extant in the Muséum national d'Histoire naturelle, Paris (MNHN). The catalogue entry lists number 7978 from "Indes Orientales" credited to "Sonnerat" as "Type de la Chagrinée de Lacepède". The holotype is later alluded to by DUMÉRIL and DUMÉRIL (1851 : 23) in the account of *Cryptopus granosus* : "Indes Orientales, Sonnerat, individu du jeune âge dont il ne reste que la carapace et la tête. Type de la Chagrinée Lacép... et de la *T. granulata* Daud." The holotype (MNHN 7978), dried and mounted, is a small *Lissemys* with the following characteristics : Skin folds cover neck to level just behind eyes ; no pattern evident on head-neck or carapace ; isolated bony prenuchal ; marginal ridge on carapace ; bony, sculptured earapace with two neurals between first pair of costals, eight pair of costals, and seven neurals (seventh indistinct) ; earapace otherwise smooth with bony peripherals posteriorly ; no limbs or tail ; femoral flaps wrinkled ; seven plastral callosities. Maximum measurements (mm) : Length (tip of snout to posterior edge of plastron), 104 ; earapace length, 77 (straight line) or 95 (wrinkled, curvature) ; carapace width, 76 (straight line) or 96 (curvature) ; bony carapace length (excluding prenuchal), 67 (straight line) or 76 (curvature) ; bony earapace width, 50 (straight line) or 56 (curvature) ; plastron length, ca. 85 ; plastron width (between lateralmost tips of hyo-hypoplastra), ca. 77. The holotype is illustrated in Fig. 1A and 1B.

#### NOMENCLATURAL HISTORY

**HOMONYMS.** — Of minor consequence is the occurrence of two junior primary homonyms, both of which were discussed by WERMUTH (1956 : 411-412). One of these, *Testudo punctata* (in SCHOEPFF, 1792 [1792-1801]) = *Emys orbicularis* (Linnaeus, 1758), is credited to GOTZWALD and declared a nomen nudum by WERMUTH (name first appears as synonym and has never been regarded as valid name). The other junior homonym, an otherwise available name, is *Testudo punctata* Schoepff, 1792 = *Clemmys guttata* (Schneider, 1792).

**REPLACEMENT NAMES.** — Three names are considered to be replacement names for *Testudo punctata* Lacepède, 1788. WERMUTH (1956 : 415-416) previously discussed the name *Testudo granulosa* Suekow, 1798, and its relevance to LACEPÈDE'S *T. punctata*, but he did not consider *T. granulosa* to be a replacement name. SUCKOW's descriptive comments (1798 : 48) are based on those of SCHNEIDER (1789 : 22-23), which in turn are based on the original description of LACEPÈDE. SCHNEIDER did not employ any scientific name, but his remarks concerning "Die chagrinirre..." and his accompanying copy of the illustration of the holotype of *T. punctata* clearly refer to LACEPÈDE's description. WER-

MUTH's interpretation was that SUCKOW was unaware of LACEPÈDE's original description and that SUCKOW's *granulosa* is based solely on SCHNEIDER's description. *Testudo granulosa* Suckow, 1798, is here regarded as a replacement name for *Testudo punctata* Lacépède, 1788.

HARPER (1940) discussed the two works of the same name "Histoire naturelle des Reptiles" by SONNINI and LATREILLE (1801) and DAUDIN (1801); he noted that the work by the former authors was published in "September or (at the latest) October, 1801" (p. 701), and that of DAUDIN in "December, 1801" (p. 715). The names *Testudo scabra* and *Testudo granulata* employed in these works are obvious replacement names for *Testudo punctata* Lacépède, 1788. The authorship of *T. scabra* is herein employed as LATREILLE in SONNINI and LATREILLE [Code, 1964, Art. 51 (e)], since it is clear that LATREILLE is solely responsible for preparation of all the accounts of species of turtles (HARPER, 1940 : 700; Roger BOUR, *pers. comm.*). The description of *T. scabra* (1801 : 164-165) is clearly based on the holotype of *T. punctata*; the account is entitled "La Tortue chagrinée, *Testudo scabra*", and the species is recorded as "apportée des Indes orientales par Sonnerat". *Testudo scabra* Latreille in SONNINI and LATREILLE, 1801, is also unavailable since it is preoccupied by two senior homonyms : 1) *Testudo scabra* Linnaeus, 1758 [= ? *Geoemyda trijuga* (Schweigger), 1812, fide Wermuth and Mertens, 1977 : 35], and 2) *Testudo scabra* Lacépède, 1788 [= *Rhinoclemmys punctularia* (Daudin, 1801), fide FRETEY, HOOGMOED, and LESCURE, 1977 : 66]. DAUDIN immediately synonymized *T. scabra* in his account of La Tortue Chagrinée (1801 : 81-85), which he called *Testudo granulata* (p. 81, footnote). DAUDIN described LACEPÈDE's young specimen, used the same common name, cited the same type-locality, and copied his illustration. DUMÉRIL and BIBRON (1835 : 504) noted that only one specimen was involved with the two names, and DUMÉRIL and DUMÉRIL (1851 : 23) mentioned the holotype of LACEPÈDE's *T. punctata* to be the type of *T. granulata*.

STATUS OF *Trionyx coromandelicus* Geoffroy. — GEOFFROY's original description of *Trionyx coromandelicus* is brief (1809 : 16-17). The diagnostic feature is the seven plastral callosities. He also alludes to the isolated bony prenuchal and peripherals, and mentions that the tail extends beyond the edge of the carapace. The following are listed as synonyms :

"Tortue chagrinée. Lacépède.  
Testudo granosa. Schoepff.  
Tortue chagrinée. Daudin.  
\_\_\_\_\_  
Bosc.  
Testudo granulata. Shaw."

The account terminates with "Patrie. La côte de Coromandel, Sonnerat, Schoepff." GEOFFROY's descriptive comments and especially his illustration of the bony carapace of *T. coromandelicus* (1809 : Pl. 5, Fig. 1) closely resemble the holotype of *Testudo punctata* Lacépède (compare bony carapaces in Figs 1A and 1C). GEOFFROY's illustration also depicts six neurals as mentioned in the original description of *T. punctata* (six neurals is an unusual variant for *Lissemys punctata*, usually seven or eight). The medial suturing posterior to the sixth neural is indistinct, but a seventh neural seems to be present (independently corroborated by Roger BOUR, *pers. comm.*). The holotype lacks a tail as pre-




FIG. 1. — A, dorsal, and B, ventral views of holotype of *Testudo punctata* Lacepède (MNHN 7978) ; C, bony carapace of *Trionyx coromandelicus*, copied from GEOFFROY (1809 : Pl. 5, fig. 1). Note overall similarity in bony carapaces (A and C).

viously noted by LATREILLE in SONNINI and LATREILLE (1801 : 165) and DAUDIN (1801 : 82-83). GEOFFROY's mention of a tail seems to be in reference to the posterior part of the plastron that does extend beyond the carapace. Certainly GEOFFROY had access to the holotype of *T. punctata* in the MNHN and I believe he simply redescribed it and supplied a new name (as he did for three other previously described species in the same paper, *Trionyx aegyptiacus*, *Trionyx stellatus*, and *Trionyx georgicus*). There is no entry for *T. coromandelicus* in the MNHN catalogue. I regard *Trionyx coromandelicus* Geoffroy, 1809, as a replacement name for *Testudo punctata* Lacepède, 1788. GEOFFROY perhaps employed the locality of "La côte de Coromandel, Sonnerat, Schoepff" (and specific name) since it is mentioned by SCHOEPFF in the original description of *Testudo granosa* (which GEOFFROY lists as a synonym) — "... in fontibus, aliisque aquis dulcibus Coromandeliae, ..." 1801 : 131, Latin edit.; "... in Quellen und anderen süßen Wassern auf Coromandel, ..." 1801 : 153, German edit. But GEOFFROY's interposition of "Sonnerat" in the above-quoted locality for *T. coromandelicus* denotes some supposition that the holotype of *T. punctata*, collected by SONNERAT, also came from the coast of Coromandel.

#### SONNERAT'S TRAVELS

The penetrating question is the geographical provenance of the holotype of *Testudo punctata* Lacepède. The holotype was received from Pierre SONNERAT, a French traveler-naturalist (born about 1745 in Lyon, died in Paris in 1814), who documented his travels in India in two quarto volumes (1782). I am thoroughly grateful to Roger BOUR, who perused these and other volumes and transmitted to me pertinent information relating to localities visited by SONNERAT.

A concise, summarizing itinerary is lacking, but the following localities are mentioned (all in Volume I) : Côte de Coromandel — Pondichéry and Karikal (p. 22); Madras, Goude-lour, and Négapatalam (p. 23); Sadras, Paliacate (p. 24); Gingi (p. 25); and Trévicarré (p. 26). Côte de Malabar — Bombay, Goa, Cochin, and Mahé (p. 34). Surate (p. 39) — this place-name is the coastal "Surat" some 240 kilometers north of Bombay. SONNERAT's activities seem to have always been in these regions. In Volume II concerning "Objets nouveaux relatifs à l'Histoire naturelle" (p. 137-248) the more interesting mammals, birds, and plants (but no reptiles) are mentioned, and in India are associated with the above-mentioned localities — the only exception is in connection with a bird, "Le petit Merle huppé des Indes" (p. 189) as living "à Malabar, à Coromandel, au Bengale" (BOUR, *pers. comm.*). There is a terminal index where "Sonnerat y a écrit quelques lignes... sur Calcutta, Chandernagor, et le Bengale. Il existe aussi un chapitre Gange (1782, I : 273-277), par 'on-dit', relatif à la religion." (Roger BOUR, *pers. comm.*).

In a new edition of SONNERAT's travels (two volumes), published in 1806 and including some annotations by Charles SONNINI is a "extrait du rapport fait par Messieurs de Lalande et de Fougeroux de Bondarois" (p. xiii-xx) (Roger BOUR, *pers. comm.*), which provides further information on SONNERAT's travels. He made three voyages as follows :

1. From 1768 to 1771, with COMMERSON, principally to Mauritius, Réunion, and Madagascar.

2. From 1771 to 1773, "de l'Inde" to the Philippines, Moluccas, and New Guinea. Details of this voyage were published by SONNERAT (1776) and a reference collection deposited "au Cabinet du Roi." This volume was consulted by Roger BOUR, who, in an effort to pinpoint the "l'Inde" locality, wrote me that SONNERAT "parti de l'Isle de France [= Mauritius], a gagné les Séchelles, puis touché seulement Ceylan, avant de se diriger, par Nicobar et Malacca, vers les Philippines, Les Moluques, pour revenir par Timor et Rodriguez, à l'Isle de France."

3. From 1774 to 1781, "repart pour Ceylan, la côte de Malabar (Mahé, Gathes), le Golfe de Cambay (Surate), la Côte de Coromandel, la Côte de l'Est (Pégu), la presqu'île Malaise, la Chine, repasse à la Côte de Coromandel (Carnate, Tanjaour, Maduré, Pondichéry), puis gagne enfin l'Isle de France, Madagascar, le Cap de Bonne Espérance" and returned with a collection of 300 birds, 50 mammals, etc. including reptiles.

In regard to the type-locality of *Testudo punctata*, Roger BOUR (*pers. comm.*) informs me that "Indes Orientales" refers to the East Indies or Indonesia, or, in a broader sense, all of southeast Asia. The "Grandes Indes" (LACEPÈDE) correspond to continental India.

It is evident that SONNERAT's activities in the Indian subregion were centered in Ceylon and the southern part of peninsular India on the east (Coromandel) and west (Malabar) coasts, but extending as far north as Surat on the west coast. All of these localities are encompassed by the geographic range of *Lissemys punctata granosa*. There is no evidence that the holotype of *Testudo punctata* originated within the confines of the Indus or Ganges river drainages. There is a possibility that the holotype came from Pegu, Burma, and, if so, represents the taxon now known as *L. p. scutata*. The data of collection strongly indicate that the holotype of *L. p. punctata* is from peninsular India. Since SONNERAT visited "Pondichéry, Côte de Coromandel" (subsequent specimens from this locality available, MNHN 9394-95, stuffed-mounted, Leschenault, 1818), this locality is chosen as the restricted type-locality of *Testudo punctata* Lacepède = Pondicherry, South Arcot (district), Tamil Nadu (state), India. Actually, Pondicherry (or Puduchcheri) is the capital of a politically distinct Union Territory of the same name.

#### TAXONOMIC HISTORY

There is certainly no direct evidence that the holotype of *Testudo punctata* is representative of the yellow-spotted, Indo-Gangetic *Lissemys*. The small holotype of *T. punctata* lacks yellow-spotting, and pattern is not mentioned in the original description.

GRAY (1831 *a* : 19 ; 1831 *b* : 49-50 ; 1832 [1830-1835] : Pl. 63) was the first author to bestow the name *punctata* on the yellow-spotted *Lissemys*, regarding that pattern characteristic of young turtles. GRAY correctly employed *punctata* as the valid name (SCHOEPFF's *Testudo granosa* and GEOFFROY's *Trionyx coromandelicus* as synonyms), and continued to use *Emyda punctata* in the ensuing years (1844 : 46 ; 1855 : 63 ; 1864 : 98 ; 1869 : 215 ; 1870 : 117 ; 1873 *a* : 71 ; 1873 *b* : 88), except for his use of *E. granosa* in one paper (1873 *c* : 306). Association of the name *punctata* with the yellow-spotted pattern (limited text supplied by GRAY) was further entrenched by BELL's Plate 4 (1838 : Part 5) copied in SOWERBY and LEAR as Plate LV (1872).

Influential authors, however, associated the yellow-spotted pattern with SCHOEPFF's *Testudo granosa*; it is of interest that SCHOEPFF acknowledges LACEPÈDE's "*T. punctata*" in his synonymy and provides some measurements of the holotype in his comparison with *T. granosa* (1801 : 127, 130-131, Latin ed.; 1801 : 148, 152, German ed.). DUMÉRIL and BIBRON (1835 : 504), employing *Cryptopus granosus*, offer the explanation that SCHOEPFF was "... le premier auteur qui ait représenté cette espèce d'une manière reconnaissable"; these authors also mention only LACEPÈDE's "La Chagrinée" in their synonymy (p. 501). This latter fact seemingly motivated BOULENGER, in his account of *Emyda granosa* (1889 : 269), to assume that LACEPÈDE (not mentioned in synonymy) failed to use binomial nomenclature. SIEBENROCK (1909 : 591), following ANNANDALE (1906 : 203), recognized two subspecies (*Emyda granosa vittata*, but referring to the nominal subspecies only as *E. granosa*); he likewise fails to note LACEPÈDE in his synonymy, and associates the yellow-spotted pattern with *E. granosa*. Later ANNANDALE (1912 : 172) associated the yellow-spotted pattern with the nominal subspecies. SMITH (1931 : 157) then provides the authoritative source for the current concept of the taxon by associating the yellow-spotted pattern with the nominal subspecies and reverting to the valid name *punctata* (but he errs in crediting the name to BONNATERRE instead of LACEPÈDE).

Thus GRAY's lead of 1831 was perpetuated by SMITH in 1931 in his concept of *Lissemys punctata punctata*. The intervening use of *granosa* as the valid name, seemingly instigated in part by LACEPÈDE's cryptic notation of binomial nomenclature, resulted in the incongruous concept of a taxon with yellow-spotted patterns correlated with a type-locality (Coromandel) where turtles having such patterns do not occur. GRAY's initial association of LACEPÈDE's *punctata* with the yellow-spotted *Lissemys* is fortuitous and unfounded.

I regard *Lissemys punctata punctata* as the valid name for the recognizable subspecies in peninsular India and Sri Lanka. The synonymy is as follows (homonyms omitted, see foregoing Nomenclatural History) :

### **Lissemys punctata punctata (Lacepède)**

*T[estudo] punctata* Lacepède, 1788 : "Synopsis methodica Quadrupedum oviparorum" [folded chart]. Type-locality, "des grandes Indes"; here restricted to Pondicherry, Coromandel Coast, India. Holotype, Mus. Nat. d'Hist. Natur. Paris 7978, stuffed-mounted young, Pierre SONNERAT, 1774-1781.

*Testudo granulosa* Suckow, 1798 : 48. Replacement name for *Testudo punctata* Lacepède, 1788.

*Testudo granosa* Schoepff, 1801 : 127 (Latin ed.); 1801 : 148 (German ed.). Type-locality, "Coromandeliae" (p. 131, Latin ed.); "auf Coromandel" (p. 153, German ed.). Two syntypes, Zool. Mus. Berlin, received by Dr. BLOCH from Dr. JOHN, not known to exist.

*Testudo scabra* Latreille in SONNINI and LATREILLE, 1801 : 164. Replacement name for *Testudo punctata* Lacepède, 1788.

*Testudo granulata* Daudin, 1801 : 81. Replacement name for *Testudo punctata* Lacepède, 1788.

*Trionyx Coromandelicus* Geoffroy St.-Hilaire, 1809 : 16. Replacement name for *Testudo punctata* Lacepède, 1788.

*Emyda vittata* Peters, 1854 : 216. Type-locality, "India orientalis, Goa." Holotype, Zool. Mus. Berlin 46, in fluid, presented by Wilhelm PETERS.

*Emyda Ceylonensis* Gray, 1855 : 64. Type-locality, "Ceylon". Type, British Museum (Natural History) 1947, 3, 4, 17, dry shell (carapace-plastron), Dr. KELAART.

[*Emyda granosa*] *intermedia* Annandale, 1912 : 172. Type-locality, "Near Purulia, Manbhumi Dist.". Holotype, Zool. Survey India 16764, in fluid.

Only ANDERSON (1876 : 514) regarded the synonymous names *Emyda granosa* and *E. punctata* as not applicable to the yellow-spotted *Lissemys*, for which he proposed the name *Emyda dura*. The name *Emyda dura*, based on a drawing of BUCHANAN-HAMILTON that has never been described or published, has been discussed elsewhere as a nomen nudum (WEBB, *in press*). Since there is no available name for the yellow-spotted, Indo-Gangetic *Lissemys*, I propose the new name :

***Lissemys punctata andersoni* new name**

HOLOTYPE. — Muséum National d'Histoire Naturelle, Paris 1977-1486, Belbari, Terai, south-eastern Nepal, elevation 210 m ; collected by Alain DUBOIS at night at the border of a pond in a forested area on 23 May 1973.




FIG. 2. — Dorsal and ventral views of holotype of *Lissemys punctata andersoni* (MNHN 1977-1486, Belbari, Nepal).

DESCRIPTION OF HOLOTYPE (Fig. 2). — Adult male (penis partly everted) ; carapace length, 189 mm ; carapace width, 145 mm ; carapace height, 63 mm ; length of plastron, 179 mm ; length of head and neck ca. 113 mm ; head width, 40 mm ; plastron without markings, showing callosities on all plastral bones (small on entoplastron) ; carapace and head with pale spots and blotches.

DIAGNOSIS AND DEFINITION. — A subspecies of *Lissemys punctata* distinguished from both *L. p. punctata* and *L. p. scutata* in having yellow markings on carapace and head.

DISTRIBUTION. — Indus and Ganges-Brahmaputra river basins in eastern Pakistan, northern India (including Sikkim and southeastern Nepal) and Bangladesh, and in north-coastal Burma. The holotype of *Lissemys punctata andersoni* represents the first record of occurrence for the taxon in Nepal.

### Acknowledgments

Although examination of study collections in various museums has contributed to the ultimate preparation of this manuscript, I am especially grateful to Dr Jean LESCURE (Muséum national d'Histoire naturelle, Paris), and to Drs T. N. Anantha KRISHNAN, Director, and Sri S. BISWAS, Zoologist (Zoological Survey of India) for the data obtained and courtesies extended. I thank The American Philosophical Society, Johnson Fund (Grant No. 1077) and The University of Texas at El Paso (Cotton Trust Funds) for partial defrayment of expenses incurred during my visit to Paris in 1977, and the Smithsonian Foreign Currency Program for providing funds to visit Calcutta in 1978. Finally, I am most grateful to Dr Roger BOUR (Mus. nat. d'Hist. natur., Paris) for reviewing the manuscript, courtesies regarding the literature, providing photographs of the holotype (Fig. 2), and for processing the manuscript for publication, without whose help this project would not have been completed. I also thank Mr Byron ALEXANDER et Roger BOUR for the drawings depicted in Fig. 1A and 1B.

### LITERATURE CITED

- ANDERSON, J., 1876. — Note on the plastron of the gangetic mud-turtle (*Enyda dura* of Buchanan-Hamilton). *J. Linn. Soc., Zool.*, **12** : 514-516.
- ANNANDALE, N., 1906. — Contributions to Oriental herpetology. No. IV. — Notes on the Indian tortoises. *J. Proc. Asiat. Soc. Beng.*, n. s., **2** (6) : 203-206, 1 pl.
- 1912. — The Indian mud-turtles (Trionychidae). *Rec. Indian Mus.*, **7** : 151-180.
- BELL, T., 1838. — A monograph of the Testudinata. Part 5. London, 5 pls.
- BONNATERRE, M., 1789. — Tableau encyclopédique et méthodique des trois règnes de la nature. Erpétologie. Paris, xxviii + 70 pp.
- BOULENGER, G. A., 1889. — Catalogue of the chelonians, rhynchocephalians, and crocodiles in the British Museum (Natural History). London, x + 311 p., 6 pls.
- DAUDIN, F. M., 1804. — Histoire naturelle, générale et particulière des reptiles... Tome Second. Paris, 1-432 p., 13 pls.
- DUMÉRIL, A. M. C., and A. H. A. DUMÉRIL, 1851. — Catalogue méthodique de la collection des reptiles du Muséum d'Histoire Naturelle de Paris. Paris, iv + 224 p.
- DUMÉRIL, A. M. C., and G. BIBRON, 1835. — Erpétologie générale ou histoire naturelle complète des reptiles. Tome Second. Paris, ii + 680 p., 12 pls.

- FRETEY, J., M. S. HOOGMOED and J. LESCURE, 1977. — Étude taxinomique de *Rhinoclemmys punctularia punctularia* (Daudin) (Testudinata, Emydidae). *Zoöl. Meded.*, **52** (6) : 63-80.
- GANS, C., 1971. — Studies on amphisbaenians (Amphisbaenia, Reptilia) 4. A review of the amphisbaenid genus *Leposternon*. *Bull. Am. Mus. nat. Hist.*, **144** (6) : 379-464.
- GEOFFROY SAINT-HILAIRE, E., 1809. — Mémoire sur les tortues molles, nouveau genre sous le nom de *Trionyx*, et sur la formation des carapaces. *Annls Mus. Hist. nat., Paris*, **14** : 1-20, 5 pls.
- GRAY, J. E., 1830-1835. — Illustrations of Indian Zoology ; chiefly selected from the collection of Major-General Hardwicke, F.R.S. Volumes I and II. London, 202 pls.
- 1831 a. — A synopsis of the species of the Class Reptilia, p. 1-110. In : E. GRIFFITH, The animal kingdom arranged in conformity with its organization, by the Baron Cuvier... London, 9 : 1-481, 33 pls.
- 1831 b. — Synopsis Reptilium ; or short descriptions of the species of reptiles. Part I, Cataphracta. Tortoises, crocodiles, and enaliosaurians. London, viii + 85 p., 12 pls.
- 1844. — Catalogue of the tortoises, crocodiles, and amphisbaenians, in the collection of the British Museum. London, viii + 60 p.
- 1855. — Catalogue of shield reptiles in the collection of the British Museum. Part I. Testudinata (Tortoises). London, 79 p., 50 pls.
- 1864. — Revision of the species of Trionychidae found in Asia and Africa, with the descriptions of some new species. *Proc. zool. Soc., Lond.*, 1864 (6) : 76-98.
- 1869. — Notes on the families and genera of tortoises (Testudinata), and on the characters afforded by the study of their skulls. *Proc. zool. Soc., Lond.*, 1869 : 165-225.
- 1870. — Supplement to the catalogue of shield reptiles in the collection of the British Museum. Part I, Testudinata (Tortoises). London, ix + 120 p.
- 1873 a. — Notes on mud-tortoises (*Trionyx*, Geoffroy), and on the skulls of the different kinds. *Proc. zool. Soc., Lond.*, 1873 : 38-72.
- 1873 b. — Hand-list of the specimens of shield reptiles in the British Museum. London, iv + 124 p.
- 1873 c. — Observations on chelonians, with descriptions of new genera and species. *Ann. Mag. nat. Hist.*, ser. 4, **11** : 289-308.
- HARPER, F., 1940. — Some works of Bartram, Daudin, Latreille, and Sonnini, and their bearing upon North American herpetological nomenclature. *Am. Midl. Nat.*, **23** (3) : 692-723.
- LACEPÈDE, B. G. E., 1788. — Histoire naturelle des quadrupèdes ovipares et des serpens. Tome Premier (quarto). Paris, 1-18, Table Méthodique..., Synopsis methodica..., 1-651 p., 41 pls.
- PETERS, W. K. H., 1854. — Hr. Peters gab eine übersicht der auf seiner reise nach Mossambique beobachteten schildkröten. *Monatsber. Akad. Wiss. Berlin*, 1854 : 215-216.
- SAVAGE, J. M., 1952. — Two centuries of confusion : The history of the snake name *Ahaetulla*. *Bull. Chicago Acad. Sci.*, **9** (11) : 203-216.
- SCHNEIDER, J. G., 1789. — Zweyter Beytrag zur Naturgeschichte der Schildkröten. Leipzig, 32 p.
- SCHOEPFF, J. D., 1792-1801. — Historia Testudinum Iconibus Illustrata (Latin ed.) ; Naturgeschichte der Schildkröten, mit Abbildungen erläutert (German ed.). Erlangen, xii + 136 (Latin ed.), xvii + 160 p. (German cd.), 34 pls.
- SHAW, G., 1802. — General zoology or systematic natural history. Vol. III, Pt. 1. London, viii + 312 p., 87 pls.
- SIEBENROCK, F., 1909. — Synopsis der rezenten Schildkröten, mit Berücksichtigung der in historischer Zeit ausgestorbenen Arten. *Zool. Jb.*, Suppl. X, **3** : 427-618.
- SMITH, M. A., 1931. — The fauna of British India, including Ceylon and Burma. Reptilia and Amphibia. Vol. I, Loricata, Testudines. London, xxviii + 185 p., 2 pls.
- SONNERAT, P., 1776. — Voyage à la Nouvelle-Guinée. Paris, xvi + 208 p., 121 pls.

- 1782. — Voyage aux Indes Orientales et à la Chine fait par ordre de Louis XVI depuis 1774 jusqu'en 1781. Paris, I : xv + 317 p., 81 pls. ; II : viii + 298 p., 61 pls.
- 1806. — Voyage... 1781 (*ibid.*) (Notes de M. Sonnini). Paris, I : xxviii + 372 + 445 p. ; II : 413 + 489 p., atlas 141 pls.

SONNINI, C. S., and P. A. LATREILLE, 1801. — Histoire naturelle des reptiles, avec figures dessinées d'après nature. Tome I, 1<sup>re</sup> partie, quadrupèdes et bipèdes ovipares. Paris, xxii + 280 p.

SOWERBY, J., and E. LEAR, 1872. — Tortoises, terrapins, and turtles drawn from life. London, iv + 16 p., I-LX pls.

SUCKOW, G. A., 1798. — Anfangsgrunde der theoretischen und angewandten Naturgeschiechte der Thiere. III. Classe. Amphibien. Leipzig, 298 p.

WEBB, R. G., *In press.* — Gray, Hardwieke, Buchanan-Hamilton, and drawings of Indian soft-shell turtles (Family Trionychidae). *Amphibia & Reptilia*, 1.

WERMUTH, H., 1956. — Versuch der deutung einiger bisher übersehener Schildkrötennamen. *Zool. Beitr.*, N. F., 2 (2-3) ; 399-423.

WERMUTH, H., and R. MERTENS, 1977. — Liste der rezenten Amphibien und Reptilien. Testudines, Croeodylia, Rhynechocephalia. Das Tierreich, Lief. 100. Berlin, xxvii + 174 p.

*Manuscrit déposé le 5 octobre 1979.*

